

CYCLES OVER FIELDS OF TRANSCENDENCE DEGREE ONE

MARK GREEN, PHILIP A. GRIFFITHS, AND KAPIL H. PARANJPE

ABSTRACT. We extend earlier examples provided by Schoen, Nori and Bloch to show that when a surface has the property that the kernel of its Albanese map is non-zero over the field of complex numbers, this kernel is non-zero over a field of transcendence degree one. This says that the conjecture of Bloch and Beilinson that this kernel is zero for varieties over number fields is precise in the sense that it is not valid for fields of transcendence degree one.

INTRODUCTION

We work over subfields k of $\mathbb{C}$, the field of complex numbers. For a smooth variety V over k , the Chow group of cycles of codimension p is defined (see [Ful98]) as

$$\mathrm{CH}^p(V) = \frac{Z^p(V)}{R^p(V)}$$

where the group of cycles $Z^p(V)$ is the free abelian group on scheme-theoretic points of V of codimension p and rational equivalence $R^p(V)$ is the subgroup generated by cycles of the form $\mathrm{div}_W(f)$ where W is a subvariety of V of codimension $(p-1)$ and f is a non-zero rational function on it. There is a natural cycle class map

$$\mathrm{cl}_p : \mathrm{CH}^p(V) \rightarrow \mathrm{H}^{2p}(V)$$

where the latter denotes the singular cohomology group $\mathrm{H}^{2p}(V(\mathbb{C}), \mathbb{Z})$ with the (mixed) Hodge structure given by Deligne (see [Del71]). The kernel of cl_p is denoted by $F^1 \mathrm{CH}^p(V)$. There is an Abel-Jacobi map (see [Gri69]),

$$\Phi_p : F^1 \mathrm{CH}^p(V) \rightarrow \mathrm{IJ}^p(\mathrm{H}^{2p-1}(V))$$

where the latter is the intermediate Jacobian of a Hodge structure, defined as follows

$$\mathrm{IJ}^p(H) = \frac{H \otimes \mathbb{C}}{F^p(H \otimes \mathbb{C}) + H}.$$

The kernel of Φ_p is denoted by $F^2 \mathrm{CH}^p(V)$.

Conjecture 1 (Bloch-Beilinson). *If V is a variety defined over a number field k then $F^2 \mathrm{CH}^p(V) = 0$.*

We (of course) offer no proof of this conjecture. However, there are examples due to C. Schoen and M. V. Nori (see [Sch87]), which show that one cannot relax the conditions in this conjecture. In this paper we present these and other examples to show that $F^2 \text{CH}^2(V)$ is non-zero for V a variety over a field of transcendence degree at least one, whenever it is non-zero over some larger algebraically closed field.

We begin in Section 1 with a lemma. We then apply this lemma to prove the following result by Hodge-theoretic methods for the second Symmetric power of a curve.

Proposition 1. *Let C be a curve of genus at least two over a number field F . There is a non-torsion cycle ξ in $F^2 \text{CH}^2(\text{Sym}^2(C) \otimes K)$, where K is the algebraic closure of the function field $F(C)$.*

In Section 2 we generalise this to surfaces other than $\text{Sym}^2(C)$. To do this we need to use an l -adic analogue of the Abel-Jacobi map. As before, let V be a smooth projective variety over k . Let R be a finitely generated subring of k for which there is a smooth projective morphism $\pi : \mathcal{V} \rightarrow \text{Spec}(R)$, whose base change to k gives V . Let $F^1 \text{CH}^p(\mathcal{V})$ denote the kernel of the natural homomorphism

$$\text{CH}^p(\mathcal{V}) \rightarrow H_{\text{ét}}^0(\text{Spec}(R), R^{2p} \pi_* \mathbb{Q}_l(p))$$

We get a natural homomorphism

$$\Phi_{p,l,\mathcal{V}} : F^1 \text{CH}^p(\mathcal{V}) \rightarrow H_{\text{ét}}^1(\text{Spec}(R), R^{2p-1} \pi_* \mathbb{Q}_l(p))$$

Since any cycle (or rational equivalence) on V over k is defined over a finitely generated ring R , the direct limit of $F^1 \text{CH}^p(\mathcal{V})$ is $F^1 \text{CH}^p(V)$. For want of better notation let us define

$$\text{IJ}_{\text{ét}}^p(H^{2p-1}(V)) = \lim_{\overline{R}} H_{\text{ét}}^1(\text{Spec}(R), R^{2p-1} \pi_* \mathbb{Q}_l(p))$$

i. e. IJ^p is the corresponding direct limit of the $H_{\text{ét}}^1$'s. An l -adic analogue of the Abel-Jacobi map is the direct limit of the $\Phi_{p,l,\mathcal{V}}$'s as we vary R ,

$$\Phi_{p,l,\text{lim}} : F^1 \text{CH}^p(V) \rightarrow \text{IJ}_{\text{ét}}^p(H^{2p-1}(V))$$

On the other hand, it is clear that we have natural homomorphisms

$$H_{\text{ét}}^1(\text{Spec}(R), R^{2p-1} \pi_* \mathbb{Q}_l(p)) \rightarrow \text{Ext}_{\text{Gal}(\overline{k}/k)}^1(\mathbb{Q}_l(-p), H_{\text{ét}}^{2p-1}(V, \mathbb{Q}_l))$$

One can show that these homomorphisms are injective when k is the field of fractions of the ring R by finding extensions of l -adic sheaves that represent elements of $H_{\text{ét}}^1$. The limiting homomorphism

$$\Phi_{p,l} : F^1 \text{CH}^p(V) \rightarrow \text{Ext}_{\text{Gal}(\overline{k}/k)}^1(\mathbb{Q}_l(-p), H_{\text{ét}}^{2p-1}(V, \mathbb{Q}_l))$$

is thus another variant of the Abel-Jacobi homomorphism that has the same kernel as $\Phi_{p,l,\text{lim}}$ when k is a finitely generated field. We use the former map to define $F^2 \text{CH}^2(V) = \cap_l \ker \Phi_{p,l,\text{lim}}$. Recent results of Nori (unpublished) show that the Hodge-theoretic definition of F^2 gives a group that is not

smaller than that given by the l -adic one above. It is still not clear that the two definitions coincide as has been conjectured by many (see [Jan95]), and has been proved by Raskind [Ras95] for *zero cycles* by using the identification of the Intermediate Jacobian with the Albanese variety in this case.

The following theorem uses the l -adic version of the definition of F^2 . In the rest of the paper the two definitions of F^2 will be used without a change of notation; it will be clear from the context which definition is being employed. By appealing to the result of Raskind [Ras95] the two definitions coincide in our case.

Theorem 2. *Let S be any surface with $p_g(S) \neq 0$ defined over a number field F . There are non-trivial cycles in $F^2 \text{CH}^2(S \otimes K, \mathbb{Q})$ where K is the algebraic closure of the function field $F(T)$ in one variable T over F .*

Acknowledgments. Kapil Paranjape would like to thank Andreas Rosenschon for discussions in which this problem was first raised. He would also like to thank M. V. Nori and S. Bloch for discussions regarding the l -adic case. We thank W. Raskind for his careful reading of the paper and numerous suggestions and corrections, especially regarding the use of limits in étale cohomology.

1. THE SECOND SYMMETRIC POWER OF A CURVE

For a surface S , let $H^2(S)_{\text{tr}}$ denote the orthogonal complement in $H^2(S)$ of the (Néron-Severi) subspace generated by cohomology classes of curves in S which are defined over an algebraically closed field.

Lemma 3 (Green-Griffiths). *Let C be a smooth projective curve and S be a smooth projective surface. Suppose Z is a homologically trivial cycle of codimension two in $C \times S$. Assume that C , S and Z are defined over a subfield k of $\mathbb{C}$. Suppose that either one of the following conditions hold:*

- (1) *The component in $\text{IJ}^2(H^1(C) \otimes H^2(S)_{\text{tr}})$ of the Abel-Jacobi invariant of Z is non-torsion. Or,*
- (2) *The component in $\text{IJ}_{\text{ét}}^2(H^1(C) \otimes H^2(S)_{\text{tr}})$ of the l -adic Abel-Jacobi invariant $\Phi_{2,l,\text{lim}}(Z)$ is non-torsion.*

Then the following map is non-torsion on $F^1 \text{CH}^1(C \otimes \mathbb{C})$,

$$z = p_{2*}(p_1^*(-) \cdot Z) : \text{CH}^1(C \otimes \mathbb{C}) \rightarrow \text{CH}^2(S \otimes \mathbb{C})$$

Proof. Let A denote the canonical divisor of C (which is defined over k). Let g denote the generic point of C considered as a point of $C \otimes k(C)$ and hence as a point of $C \otimes \mathbb{C}$ via some inclusion of $k(C)$ in $\mathbb{C}$. If $z(\deg(A)g - A)$ is torsion in $\text{CH}^2(S \otimes \mathbb{C})$, then a rational equivalence demonstrating this is defined over some finite extension of $k(C)$. Taking the norm of this equivalence we see that $z(\deg(A)g - A)$ is torsion in $\text{CH}^2(S \otimes k(C))$ as well. Hence a multiple of $z(\deg(A)g - A)$ is a sum of the form $\sum \text{div}_{\mathcal{D}_i}(f_i)$ where $\mathcal{D}_i \subset \text{Spec}(k(C)) \times S$ is a curve and f_i is a function on $\mathcal{D}_i$. Let $D_i \subset C \times S$ be the Zariski closure of $\mathcal{D}_i$ and consider the cycle W obtained by summing up $\text{div}_{D_i}(f_i)$. By

assumption, W restricts to a multiple of $z(\deg(A)q - A)$ on $\text{Spec}(k(C)) \times S$. Thus there are closed points q_j of C over k and divisors E_j in $\text{Spec}(k(q_j)) \times S$ so that for some integer n we have

$$\sum \text{div}_{D_i}(f_i) = W = n(\deg(A)Z - C \times z(A)) + \sum_j q_j \times E_j$$

The image of cycles of the form $C \times z(A)$ as well as cycles of the form $\sum_j q_j \times E_j$ in the relevant piece of IJ^2 is zero in case (1) as well as (2). This contradicts the hypothesis on Z . $\square$

From this lemma it follows that in order to construct a non-trivial cycle in $F^2 \text{CH}^2(S)$ over a field of transcendence degree one it is sufficient to construct C , S and Z as above over a field of transcendence degree 0.

Corollary 4. *Let C be a smooth projective curve and S a smooth projective surface over a subfield F of $\overline{\mathbb{Q}}$. Let Z be a homologically trivial cycle on $C \times S$ such that the induced homomorphism*

$$z = p_{2*}(p_1^*(-) \cdot Z) : \text{CH}^1(C \otimes \mathbb{C}) \rightarrow \text{CH}^2(S \otimes \mathbb{C})$$

is non-torsion. Then, there is a non-torsion class in $F^2 \text{CH}^2(S \otimes_F \overline{F(C)})$.

Proof. From a result of Roitman [Roj80] it follows that the kernel of z is a countable union of translates of proper abelian subvarieties A_i of $J(C)$. But each such translated abelian subvariety is defined over $\overline{\mathbb{Q}}$ since Z is defined over $\overline{\mathbb{Q}}$. Using a point p of C over $\overline{\mathbb{Q}}$ we embed C in $J(C)$. Since the image of C generates $J(C)$, the image of the generic point $\text{Spec}(F(C)) \rightarrow C$ is not in A_i for any i . Its image under z gives us the desired cycle in $F^2 \text{CH}^2(S \otimes_F \overline{F(C)})$. $\square$

Let C be a curve of genus at least 2 over $k = \overline{\mathbb{Q}}$. Let $S = \text{Sym}^2(C)$ and consider the cycle X in $C \times C \times S$ which is the graph of the natural map $q : C \times C \rightarrow S = \text{Sym}^2(C)$. The map q_* on $H^{1,0}(C) \otimes H^{1,0}(C)$ identifies the range $H^{2,0}(S)$ with the second exterior power of $H^{1,0}(C)$. The cohomology class $[X]$ therefore induces a non-zero map

$$p_{23*}(p_1^*(-) \cup [X]) : H^{1,0}(C) \rightarrow H^{0,1}(C) \otimes H^{2,0}(S)$$

Thus the map

$$J(C \otimes \mathbb{C}) = \text{IJ}^1(H^1(C)) \rightarrow \text{IJ}^2(H^1(C) \otimes H^2(S)_{\text{tr}})$$

is non-torsion. Hence, the image is a non-zero abelian variety A which is a quotient of $J(C \otimes \mathbb{C})$. Now, $J(C)$ is defined over $\overline{\mathbb{Q}}$ and hence so is any quotient abelian variety. In particular, A is $B \otimes \mathbb{C}$ for some abelian variety B defined over a number field.

Proposition 5. *Let $J \rightarrow B$ be a surjective homomorphism of abelian varieties over a number field; $B \neq 0$. There are points in $J(\overline{\mathbb{Q}})$ whose image in B is non-torsion.*

Proof. Let L be the completion of the number field at a place lying over a prime p and M be the algebraic closure of $\mathbb{Q}$ in L . $J(L)$ and $B(L)$ are p -adic Lie groups. Consider a finite morphism $\pi : J \rightarrow \mathbb{P}^n$ and let $U \subset \mathbb{P}^n$ be a Zariski open set over which π is étale. It is clear that $U(M)$ is dense in $U(L)$. By the implicit function theorem the map $J(L) \cap \pi^{-1}(U) \rightarrow U(L)$ is open; moreover, the former set is dense in $J(L)$. Since π is finite, any point in $J(L) \cap \pi^{-1}(U(M))$ is in $J(M)$. Thus $J(M)$ is dense in $J(L)$. Now $J(L) \rightarrow B(L)$ is open, so the image of $J(M)$ in $B(L)$ is dense in an open subgroup. The torsion subgroup of a p -adic Lie group is finite. The result follows. $\square$

Corollary 6. *Let A be a non-zero abelian variety over a number fields. The group $A(\overline{\mathbb{Q}})$ is non-torsion.*

By this proposition and the discussion above we obtain a cycle Z on $C \times S$ over $\overline{\mathbb{Q}}$ as required. Now applying the lemma we see that we have proved Proposition 1.

2. THE GENERAL CASE

We use an argument of Terasoma (as explained to us by Bloch and Nori) to construct a triple (C, S, Z) as in Lemma 3 for *any* surface S with $p_g \neq 0$.

Lemma 7. *Assume $(\tilde{C}, \tilde{S}, \tilde{Z})$ is a triple over some finitely generated subfield k of $\mathbb{C}$ which satisfies the condition (2) of Lemma 3. Then, there exists a specialisation (C, S, Z) over a number field $F \subset \overline{\mathbb{Q}}$ which satisfies the condition (2) of Lemma 3.*

Proof. Let T be an affine variety over a number field contained in k and (C, S, Ξ) be a triple over S so chosen that this data restricts to $(\tilde{C}, \tilde{S}, \tilde{Z})$ under the natural map $\text{Spec } k \rightarrow T$. Using the Gysin sequence, we obtain an exact sequence of $\mathbb{Q}_l$ constructible local systems on T ,

$$0 \rightarrow R_{\text{ét}}^1 \pi_{T*}(\mathcal{C}, \mathbb{Q}_l) \otimes R_{\text{ét}}^2 \pi_{T*}(\mathcal{S}, \mathbb{Q}_l)_{\text{tr}} \rightarrow \mathcal{V} \rightarrow \mathbb{Q}_l(-2)_T \rightarrow 0$$

here $R_{\text{ét}}^2 \pi_{T*}(\mathcal{S}, \mathbb{Q}_l)_{\text{tr}}$ denotes the local system on T associated with the Galois representation $H_{\text{ét}}^2(\tilde{S} \otimes \overline{k}, \mathbb{Q}_l)_{\text{tr}}$. Moreover, the extension class of this short exact sequence is naturally identified with the component in $H_{\text{ét}}^1(T, R_{\text{ét}}^1 \pi_{T*}(\mathcal{C}, \mathbb{Q}_l) \otimes R_{\text{ét}}^2 \pi_{T*}(\mathcal{S}, \mathbb{Q}_l)_{\text{tr}} \otimes \mathbb{Q}_l(2))$ of the class $\Phi_{2,l,C \times_T S}(\Xi)$. By assumption, condition (2) of Lemma 3 means this component is non-zero. So the sequence is non-split.

If p is a closed point of T , let C_p, S_p, Z_p denote the specialisations. By restricting the above sequence we obtain an exact sequence of representations of $\text{Gal}(\overline{\mathbb{Q}}/k(p))$ or equivalently, of a decomposition group of p in $\text{Gal}(\overline{k}/k)$,

$$0 \rightarrow H_{\text{ét}}^1(C_p \otimes \overline{\mathbb{Q}}, \mathbb{Q}_l) \otimes (R_{\text{ét}}^2 \pi_{T*}(\mathcal{S}, \mathbb{Q}_l)_{\text{tr}})_{\overline{k(p)}} \rightarrow \mathcal{V}_{\overline{k(p)}} \rightarrow \mathbb{Q}_l(-2) \rightarrow 0$$

There are two possible problems.

- (1) There may be more divisors in S_p than on the generic fibre of $\mathcal{S} \rightarrow T$ so that

$$H_{\text{ét}}^2(S_p \otimes \overline{k(p)}, \mathbb{Q}_l)_{\text{tr}} \text{ is a proper subset of } (R_{\text{ét}}^2 \pi_{T*}(\mathcal{S}, \mathbb{Q}_l)_{\text{tr}})_{\overline{k(p)}}$$

- (2) The above exact sequence may split so that the extension class becomes zero for Z_p .

These two problems are resolved by applying the following lemma to the l -adic representation

$$\mathcal{V}_{\overline{k}} \oplus H_{\text{ét}}^2(S \otimes \overline{k}, \mathbb{Q}_l)$$

□

Lemma 8 (Terasoma). *Let V be a continuous l -adic representation of the fundamental group of a variety T over a number field. Then there are infinitely many closed points p of T so that the image of $\text{Gal}(\overline{k(T)}/k(T))$ in $\text{GL}(V)$ is the same as the image of the decomposition group of p .*

Proof. Let G denote the image of $\text{Gal}(\overline{k(T)}/k(T))$ in $\text{GL}(V)$ and D_p denote the image of a decomposition group associated with a closed point p of T . These are compact subgroups of $\text{GL}(V)$. As in [Ter85], there is a subgroup H of finite index in G so that $D_p = G$ if and only if the natural map $D_p \rightarrow G/H$ is a surjection. Equivalently, if $f : T' \rightarrow T$ denotes the étale cover of T' corresponding to this finite quotient of the fundamental group of T , then it is enough that $f^{-1}(p)$ is a closed point of T' . To find such a point p , we may shrink T to assume that T is affine and that there is an étale map $g : T \rightarrow \mathbb{A}^n$. Thus it is enough to find a point q in $\mathbb{A}^n$ so that $(g \circ f)^{-1}(q)$ is a closed point of T' . Such a point can be found by the Hilbert irreducibility theorem. □

Now we consider a surface S over $\overline{\mathbb{Q}}$ with $p_g(S) \neq 0$; let p be a closed point on S . Murre (see [Mur90]) has constructed a decomposition of the diagonal of a surface S ,

$$\Delta_S = p \times S + S \times p + X_{2,2} + X_{1,3} + X_{3,1}$$

for the diagonal Δ_S in $\text{CH}^2(S \times S) \otimes \mathbb{Q}$; where $X_{i,j}$ is a cycle class so that its cohomology class has a non-zero Künneth component only in $H^i(S) \otimes H^j(S) \otimes \mathbb{Q}$. Moreover, by the Hodge index theorem we can write $nX_{2,2} = \sum_i C_i \times D_i + X_{2,2,\text{tr}}$ where C_i and D_i are curves on S and $X_{2,2,\text{tr}}$ has its cohomology class in $H^2(S)_{\text{tr}} \otimes H^2(S)_{\text{tr}}$.

Let $H \subset S \times \mathbb{P}^1$ be any pencil; $Y_{i,j}$ denote the cycles on $H \times S$ that are the natural pull-backs of $X_{i,j}$ for each i and j ; similarly, let $Y_{2,2,\text{tr}}$ be the pull-back of $X_{2,2,\text{tr}}$. Let H_0 denote the divisor class of the fibres of the pencil and A the class of an ample divisor on S . The class $Y_{2,2,\text{tr}}$ on $H \times S$, restricts to a homologically trivial class on every member H_t of the pencil; however, $H^2(H)_{\text{tr}} = H^2(S)_{\text{tr}}$ so that the cohomology class of $Y_{2,2,\text{tr}}$ in $H^2(H)_{\text{tr}} \otimes H^2(S)_{\text{tr}}$ is non-zero.

For any open subset $U \subset \mathbb{P}^1$, let $H_U = H \times_{\mathbb{P}^1} U$; the map $H^2(H)_{\text{tr}} \rightarrow H^2(H_U)$ is injective. Thus the pull-back of $Y_{2,2,\text{tr}}$ to $H_U \times S$ is not homologically trivial for all U . By the Leray spectral sequence for the map $\pi : H_U \rightarrow U$, we see that $H^2(H)_{\text{tr}}$ maps injectively into $H^1(U, R^1 \pi_*(\mathbb{Q}_l))$. Let g denote the generic point of $\mathbb{P}^1$. The restriction of $Y_{2,2,\text{tr}}$ to $H_g \times S$ is a cycle Z_g such that the triple $(H_g, S \otimes_k k(g), Z_g)$ satisfies the condition (2) of Lemma 7 over the field $k(g)$. Applying Lemma 7 we see that we have a point t in $\mathbb{P}^1(\mathbb{Q})$ so that (H_t, S, Z_t) is a triple satisfying condition 3 of the Lemma 3 of Green and Griffiths. Thus we have proved Theorem 2.

REFERENCES

- [Del71] Pierre Deligne, *Théorie de Hodge. II*, Inst. Hautes Études Sci. Publ. Math. (1971), no. 40, 5–57.
- [Ful98] William Fulton, *Intersection theory*, second ed., Springer-Verlag, Berlin, 1998.
- [Gri69] Phillip A. Griffiths, *On the periods of certain rational integrals. I, II*, Ann. of Math. (2) **90** (1969), 460–495; *ibid.* (2) **90** (1969), 496–541.
- [Jan95] Uwe Jannsen, *Mixed motives, motivic cohomology, and Ext-groups*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 667–679.
- [Mur90] J. P. Murre, *On the motive of an algebraic surface*, J. Reine Angew. Math. **409** (1990), 190–204.
- [Ras95] Wayne Raskind, *Higher l -adic Abel-Jacobi mappings and filtrations on Chow groups*, Duke Math. J. **78** (1995), no. 1, 33–57.
- [Roj80] A. A. Rojtman, *The torsion of the group of 0-cycles modulo rational equivalence*, Ann. of Math. (2) **111** (1980), no. 3, 553–569.
- [Sch87] Chad Schoen, *Zero cycles modulo rational equivalence for some varieties over fields of transcendence degree one*, Algebraic Geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), Amer. Math. Soc., Providence, RI, 1987, pp. 463–473.
- [Ter85] Tomohide Terasoma, *Complete intersections with middle Picard number 1 defined over $\mathbb{Q}$* , Math. Z. **189** (1985), no. 2, 289–296.

DEPT. OF MATH., UCLA, LOS ANGELES, CALIFORNIA, USA.
E-mail address: `mlg@ipam.ucla.edu`

IAS, PRINCETON, NEW JERSEY, USA.

IMSC, CIT CAMPUS, THARAMANI, CHENNAI 600 113, INDIA.
E-mail address: `kapil@imsc.res.in`